

# 0-0-13 High K Fertilizer Plus Preemergent Weed Control



## FEATURES & USE TIPS:

- Combination fertilizer and Pendimethalin preemergent herbicide.

- Provides unsurpassed preemergent control of:

### **Annual Grassy Weeds:**

Crabgrass	Foxtail	Fall panicum
Goosegrass	Barnyardgrass	<i>Poa annua</i>

### **Broadleaf Weeds (before they germinate):**

Chickweed	Knotweed	Hop clover
Oxalis	Purslane	Henbit
Prostrate Spurge	Cudweed	Eveningprimrose

- Consistently sized particles reduce segregation potential
- Small particle formulation more particles per square inch for better weed control.
- Special no-nitrogen formula

## PRODUCT SPECIFICATIONS:

**K - as K<sub>2</sub>O - (13%):** Potassium chloride

**Cl (Not more than 13%):** Potassium chloride

**Active Ingredient:** Pendimethalin (0.84%)

**PRODUCT CODE:**

**AGC86013**

Date: 101002  
Supersedes: 000000

## PRODUCT CHARACTERISTICS:

### **Product Type:**

Combination fertilizer and Pendimethalin pre-emergent herbicide

### **Particle Size:**

SGN: 150

### **Net Weight:**

50 lbs.

### **Coverage:**

10,700 sq. ft. at the medium rate

Andersons Golf Products is a trademark of The Andersons Agriservices, Inc.

*To order, contact your local Andersons Golf Products distributor.*

*For more information, contact Andersons at our website: [www.andersonsgolfproducts.com](http://www.andersonsgolfproducts.com) or call 1-800-225-2639.*



**ACTIVE INGREDIENT**

Pendimethalin†	0.84%
<b>OTHER INGREDIENTS</b>	<b>99.16%</b>
Total	100.00%

†CAS No. 40487-42-1

EPA Reg. No. 9198-209

EPA Est. No. 538-OH-1S, 9198-OH-1M

Underlined letter is first letter used in run code on bag.

**GUARANTEED ANALYSIS**

Soluble Potash (K <sub>2</sub> O)	13.00%
Plant nutrients derived from potassium chloride.	
Chlorine (Cl) Max.	13.00% <b>F292</b>

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-ap.com>.

U.S. Pat. Nos. 5,219,465; 5,300,135; 5,405,426 and 5,466,274

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals:** CAUTION, causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed.

**FIRST AID**

**If swallowed:** Call a poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

**Personal Protective Equipment**

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves such as butyl rubber, barrier laminate, or nitrile rubber greater than 14 mils (if you want more options, follow instructions for Category A on EPA Chemical-Resistant Selection Chart), shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining personal protection equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash personal protection equipment separately from other laundry.

**User Safety Recommendations:**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**Environmental Hazards:** This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high

water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

Prevents sprouting of the following grassy and dicot weeds and provides a feeding of potassium:

**Grassy Weeds**

- Barnyardgrass
- Crabgrass
- Fall panicum
- Foxtail
- Goosegrass
- Poa annua*

**Dicot Weeds**

- Chickweed
- Cudweed
- Eveningprimrose
- Knotweed
- Henbit
- Hop clover
- Oxalis
- Prostrate spurge
- Purslane

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves such as butyl rubber, barrier laminate, or nitrile rubber greater than 14 mils (if you want more options, follow instructions for Category A on EPA Chemical-Resistant Selection Chart), shoes plus socks.

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, tall fescue, fine fescue, bentgrass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, centipedegrass, bahiagrass, and zoysiagrass and blends of the grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions" and "Limitation of Liability" Statements.

Not for use on bentgrass, *Poa annua*, and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

**How/When to Use**

For control of annual grassy and broadleaf weeds such as crabgrass, goosegrass, foxtail, barnyardgrass, fall panicum, *Poa annua*, oxalis, knotweed, purslane, and prostrate spurge, apply in the spring before they sprout.

To prevent *Poa annua* and fall germinating chickweed, apply product to dry foliage in late summer-early fall (before germination). Repeat application yearly

**0-0-13 High K Fertilizer  
Plus Preemergent Weed Control**



for continued control. A repeat application made after *Poa annua* treatment should not be sooner than 4 months.

Where lower infestations are expected, lighter rates may be used.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after applications (1/4 - 1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying as a segment to application. Allow at least a 4 month span between the applications except when repeat applications are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.

**Precautions**

Do not use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's. This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

Do not apply to dormant bentgrass and *Poa annua* (turf that is not actively growing or has experienced frost).

When any preemergence herbicide is applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring. The delay in green-up may be extensive in low-cut bermudagrass and on sandy soils.

As with most other preemergent herbicides, application of this product to bermudagrass overseeded with cool season turfgrasses may injure and thin the overseeding.

Do not apply to bentgrass and *Poa annua* after late summer.

Avoid application to cement, stone, clothing, or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 mph) due to its staining potential.

Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

**Properties of All Dinitroaniline (DNA) Products**

May cause temporary slowing of turf fill-in and stolon-rhizome rooting where turf has been thinned.

Requires a waiting period before seeding or sprigging.

**Seeding or Sprigging Intervals after Treatment**

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

**Treatment Interval after Seeding or Sprigging**

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

**Notes**

Do not apply more than a cumulative total of 349 lbs. product/acre (3.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua*; 698 lbs. product/acre (6.0 lbs. active ingredient/acre) per year on cool season turfgrass and 1,046 lbs. product/acre (9.0 lbs. active ingredient/acre) per year on warm season turfgrass.

TURFGRASS SPECIES	WEEDS	RATES OF TURF FERTILIZER PLUS PREEMERGENT WEED CONTROL	COMMENTS
<b>• Cool Season Grasses</b> Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	Crabgrass Foxtail <i>Poa annua</i> Barnyardgrass Fall panicum Oxalis Prostrate spurge Purslane Knotweed Chickweed	175-233 lbs./acre as the initial application prior to weed germination.	Apply a repeat application of 145-175 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	Goosegrass	175-233 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6-8 weeks at 175-203 lbs./acre.	
Bentgrass and established <i>Poa annua</i> (1/2 inch height or taller. Not for use on greens and tees).	Crabgrass Foxtail <i>Poa annua</i> Barnyardgrass Fall panicum Oxalis Prostrate spurge Purslane Knotweed Chickweed	175 lbs./acre in the spring prior to weed germination.	Apply a repeat application of 145-175 lbs./acre after 6-8 weeks where heavy infestations are expected.
	Goosegrass	175 lbs./acre in the spring prior to weed germination with a repeat application of 175 lbs./acre after 6-8 weeks.	
<b>• Warm Season Grasses</b> Bermudagrass St. Augustinegrass Bahiagrass Zoysiagrass Centipedegrass (401 lbs./acre rate or less) Tall fescue (Not for use on greens and tees.)	Crabgrass Foxtail <i>Poa annua</i> Barnyardgrass Fall panicum Oxalis Prostrate spurge Purslane Knotweed Chickweed	175-233 lbs./acre prior to weed germination.	Apply a repeat application of 145-175 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	Goosegrass	175-233 lbs./acre prior to weed germination	Apply a repeat application of 175-233 lbs./acre to extend control of goosegrass if the lower rate was used initially.
	Cudweed Hop clover Eveningprimrose Henbit	175-233 lbs./acre prior to weed germination.	Apply a repeat application of 145-203 lbs./acre after 6-8 weeks, if the lower rate was used initially.

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION of LIABILITY**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (APtrans/RVnot 072301)

**Suggested Spreader Settings**

These suggested settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. feeds 15,000 sq. ft. (0.34 acre) at the LIGHT RATE

50 lbs. feeds 12,400 sq. ft. (0.28 acre) at the LOW RATE

50 lbs. feeds 10,700 sq. ft. (0.25 acre) at the MEDIUM RATE

50 lbs. feeds 9,300 sq. ft. (0.21 acre) at the HIGH RATE

**SPREADER SETTINGS**

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LIGHT RATE 3.33 lbs./k sq. ft.	LOW RATE 4.02 lbs./k sq. ft.	MEDIUM RATE 4.66 lbs./k sq. ft.	HIGH RATE 5.35 lbs./k sq. ft.
Andersons Rotaries: AccuPro/SR-2000 (Cone 8)	3 mph	12 feet	J	J 1/2	K	K 1/2
Andersons SS-2	3 mph	overlap wheels	4 1/2	5	5 1/4	5 3/4
Scotts Rotaries: R-8, R-8A (Cone 8)	3 mph	12 feet	I 1/2	J	J 1/2	K 1/2
Lely Models WTR, WFR, HR, 1250. (PTO at 450 rpm)	4.5 mph	36 feet	4 3/4 I	5 I	5 1/4 I	5 1/2 I
Vicon (03 Series-Standard Spout)	4.5 mph	28 feet	24	27	29	32

Note: These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

**AGC86013**

Product of USA

**Manufactured by:**

The Andersons Lawn Fertilizer Division, Inc.  
P. O. Box 119, Maumee, OH 43537  
1-800-225-2639

Visit us on the web at: [www.AndersonsGolfProducts.com](http://www.AndersonsGolfProducts.com)