

Reinders

20-0-10

with 0.72% TRIMEC® Herbicide

• Controls dandelion, chickweed, knotweed, plantain, henbit and other broadleaf weeds as listed

• Contains polymer coated sulfur coated urea slow release nitrogen

GUARANTEED ANALYSIS:

Total Nitrogen (N)	20.00%
20.00% Urea Nitrogen*	
Soluble Potash (K ₂ O)	10.00%
Sulfur (S)	1.80%
1.80% Free Sulfur (S)	
Plant nutrients derived from urea, potassium chloride and polymer coated sulfur coated urea.	
Chlorine (Cl) Max	8.50%
*6.00% Slowly available nitrogen from polymer coated sulfur coated urea.	F292

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid	0.55%
(+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	0.12%
Dicamba: (3,6-dichloro-o-anisic acid)	0.05%
INERT INGREDIENTS	99.28%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

11.0 lbs. 2,4-Dichlorophenoxyacetic acid equivalent per ton
 2.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton
 1.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton
 Isomer Specific by AOAC Methods.
 Contains the single isomer form of Mecoprop-p.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Below For Additional Precautionary Statements

EPA Reg. No.2217-581-9198
 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
 Underlined letter is first letter used in run code on bag.

**STOP! READ THE ENTIRE LABEL FIRST.
 OBSERVE ALL PRECAUTIONS AND FOLLOW
 DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

Caution: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product wear long-sleeved shirt, long pants, gloves, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

Trimec® is a registered trademark of PBI/GORDON Corp.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free 1-877-800-5556 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from target area.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPP-p and 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPP-p and 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. **DISPOSAL: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WHERE TO USE

This product is a TRIMEC® brand herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

This product is intended to be applied by noncommercial users, lawn care operators and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example hospitals, nursing homes, schools, museums, libraries, sport

facilities (including golf courses) and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

This product may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system. This product may be applied to newly established warm season turfgrass. Treatments should be delayed until 3-4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the recommendation for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Do not use this product on dichondra, carpetgrass, St. Austinegrass, closely mowed bentgrass putting greens, or turf containing desirable clovers.

WEEDS CONTROLLED

Dandelion, chickweed, plantain, and clover are controlled effectively with this product. A partial list of susceptible weeds is shown below:

BEDSTRAW	DOCK	PIGWEED
BLACK MEDIC	GROUND IVY	PLANTAIN
BUCKHORN	HEALALL	PURSLANE
BURDOCK	HENBIT	RAGWEED
CHICORY	KNOTWEED	SHEEP SORREL
CHICKWEED	LAMBSQUARTERS	SHEPHERDSPURSE
CLOVER	MORNINGGLORY	SPEEDWELL
DANDELION	PEPPERGRASS	THISTLE

HOW MUCH TO APPLY

FOR KENTUCKY BLUEGRASS, FESCUE spp., PERENNIAL RYEGRASS, BERMUDAGRASS, AND ZOYSIAGRASS: Apply 139 - 218 lbs. of this product per acre of established turfgrass. Or, the equivalent application rate is 3.2 - 5.0 lbs. per 1,000 sq. ft. of turfgrass.

Generally the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

The maximum application rate to turf is 1.2 lbs. 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer.

Refer to the rate table for the appropriate bag sizes and the area to be treated.

DESIRED RATES		TREATED AREA FOR THIS BAG SIZE	
Per Acre, Pounds	Per 1,000 Sq. Ft., Pounds	Acres	Sq. Ft.
139 - 218	3.2 - 5.0	0.23 - 0.36	10,000 - 15,600

Application Suggestions:

- Spring and fall applications are recommended for best control. However, this product can be applied any time broadleaf weeds are present and actively growing.
- Mow the turfgrass 2 to 3 days before application.
- Water the turfgrass thoroughly (> 1/2 inch) at least 1 to 2 days before application.

- Schedule applications in the morning when dew is present. Or sprinkle or irrigate lightly (< 1/4 inch) before application to provide maximum control.
- Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

Suggested Spreader Settings				
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.				
50 lbs. covers 15,600 sq. ft. (0.36 acre) with 0.64 lbs. N/1,000 sq. ft. at the LOW RATE				
50 lbs. covers 12,500 sq. ft. (0.29 acre) with 0.80 lbs. N/1,000 sq. ft. at the MEDIUM RATE				
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LOW RATE 3.2 lbs/1,000 sq. ft.	MEDIUM RATE 4.0 lbs/1,000 sq. ft.
Andersons AccuPro 2000, SR-2000 (cone 4)	3 mph	12 feet	K 3/4	L 1/2
Andersons AccuPro 1000	3 mph	12 feet	K 3/4	L 1/2
Scotts R-8, R-8A (cone 4)	3 mph	12 feet	J 1/4	K
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern II-A	4.5 mph	34 feet	4	4 3/4
Vicon (O3 Series)	4.5 mph	32 feet	20	24

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of USA

Distributed by:
 Free Flow Fertilizer
 PO Box 119
 Maumee, OH 43537

SGN 215
Net Wt. 50 Lbs. (22.68 kg)

RND206WF5

B8 (BLKP1929B)