

[Click here to print this page](#)[Close Window](#)

Specimen Label

Neutralize

Spray Tank Systems Cleaner

Non-Corrosive

Helps prevent cross-contamination

One pound treats 100 gallons

Safe for Agricultural, Turf and Pest Control use

DIRECTIONS FOR USE

Fill spray tank to half capacity with water.

Engage agitation.

Slowly add 1 pound (454 g) of Neutralize for every 100 gallons (379 ltr) of water added to the tank.

Agitate throughout tank for approximately 15 minutes.

Rinse all internal areas of the tank, then empty the tank by the boom, and other outlets to ensure all are clean.

Close valves and refill tank with clean water. Empty water through all outlets and boom to ensure no residue remains.

Test the second rinse solution (See Method #1 or #2 under testing) to ensure no pesticide residue remains. If pesticide residue remains in the solution, repeat the cleaning process.

Exact rates should be determined by the user as rates will vary depending on the type, size, and condition of the spray tank.

BACKPACK USE

Add 1/2 ounce (14 g) of Neutralize (1 full tablespoon) to a 3 gallon (11 ltr) backpack. Fill sprayer with water and shake. Empty the rinsate through the hand wand. Be sure all pesticide is removed prior to reusing.

SPECIAL INSTRUCTIONS

Rubber parts, such as hoses, pump packing, etc. may retain pesticide residues. Exterior areas of spray tanks may be cleaned with Neutralize.

Neutralize can prevent corrosion and dissolution of metals. All rinsate must be properly handled since it may contain pesticides residue.

RINSATE TESTING

Certain pesticides, including but not limited to 2,4-Ds and Sulfonylureas, are difficult to remove from the tank. Pesticides are extremely active at very low rates and must be completely removed from the tank to prevent crop damage. Do not apply Neutralize or rinsate directly to crops. Rinsate levels may be tested by several different methods; two are suggested below:

Method One: Retain a sample of the rinsate solution. Apply sample to sensitive plants such as an ornamental flower, tomato, or cotton. Observe treated plant for 48 hours or longer, and do not use tank during the test period. If plant damage occurs, reclean the tank until the rinsate is pesticide

free.

Method Two: Retain a sample of rinsate solution and submit to a qualified testing laboratory for analysis. Operators of multiple chemical use equipment and growers of high value or sensitive crops may wish to use all available testing methods to prevent accidental contamination and crop damage.

Warning

Contains sodium carbonate, CAS Registry No. 497-19-8. Causes eye and skin irritation. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. In case of contact, immediately flush eyes and skin with plenty of water for at least 15 minutes. Call a physician. In case of spillage, sweep, scoop or vacuum and remove. Flush spill area with water. For additional information, see Material Safety Data Sheet (MSDS). For Industrial Use Only.

KEEP OUT OF REACH OF CHILDREN

EMERGENCY CONTACT CHEMTREC: 1-800-424-9300

NOTICE: There are no warranties which extend beyond the description on the label hereof. The manufacturer, Becker Underwood, Inc., warrants this product to be as described on the label. The manufacturer's liability on this warranty is limited to return of the purchase price of this product. The buyer assumes all risks and liability resulting from the use of this product, whether used singly or in combination with other goods. The manufacturer neither assumes nor authorizes any person to assume for it any other liability in connection with the sale or use of this product.

Manufactured By:

Becker Underwood, Inc.

801 Dayton Avenue
Ames, IA 50010
515-232-5907
Fax 515-232-5961
www.bucolor.com

Net Weight: 32 oz. (907 grams)

[Click here to print this page](#)

[Close Window](#)