



26/36[®]

FUNGICIDE

Dual Mode of Action for Effective Disease Control.

Incorporating 26/36[®] fungicide into an integrated disease management program provides effective control of key diseases of turfgrass and ornamentals.

EASY-TO-USE FORMULA

- Easy-to-use, premixed formulation
- Convenient
- Broad-spectrum disease control for turf
- Tank-mix compatible with most commonly used pesticides

EPA REG. NO. 228-630

ACTIVE INGR. iprodione (19.65%),
thiophanate-methyl (19.65%)

FORMULATION flowable

CHEM. FAMILY dicarboxamide, benzimidazole

FRAC NO. 1 and 2

SIGNAL WORD warning

RESTRICTED no

PACKAGE SIZE 2 x 2.5 gal

 COMMERCIAL

 TURF

 LANDSCAPE ORNAMENTALS

 GOLF TURF



AN EXCLUSIVE LINE FROM NUFARM



26/36[®]

FUNGICIDE

KEY USES

Golf courses
Institutional areas
Ornamentals
Sod farms
Turfgrass

KEY DISEASES CONTROLLED

(See product label for complete list)

Anthracnose
Ascochyta leaf blight
Brown patch
Copper spot
Corticium
Dollar spot
Fusarium patch
Gray snow mold
Leaf spot
Pink snow mold
Red thread
Zoysia patch

APPLICATION RECOMMENDATIONS

26/36[®] fungicide is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, and institutional areas where fine turf is grown. It provides effective, broad spectrum turf disease control and also serves as a useful tank mixture in the resistance management program required for other resistant sensitive fungicides.

When used in conjunction with good turf management practices, 26/36 Fungicide is effective in controlling the following diseases: dollar spot, brown patch, anthracnose, leaf spots such as Helminthosporium leaf spot caused by Drechslera spp. pathogens, corticium red thread, Fusarium patch, zoysia patch, ascochyta leaf blight and copper spot.

Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

See label for complete application rates and recommendations.

PRECAUTIONS AND RESTRICTIONS

Do not exceed a total of 14.5 fl oz product per 1000 sq ft per year.

Under severe conditions, the higher rate and/ or shorter interval of applications are recommended for all diseases. When disease pressure is low to moderate, the lower rates and longer intervals are recommended.

Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24-hour waiting period following treatment is preferred.

Do not mix with any sticker, extender, or wetting agent.

Do not graze animals on treated turf.

Do not feed clippings from treated turf to livestock or poultry.

See label for complete application precautions and restrictions.



More where it matters most.
www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.
©2013 Nufarm. Important: Always read and follow label instructions. 26/36[®] is a trademark of Cleary Chemical, LLC. 13-TO-0069



Scan this code with your smartphone to visit CDMS and view the complete product label.