

Active Ingredients: fenoxaprop-p-ethyl **Formulation:** suspension emulsion

WSSA Group: 1 Signal Word: caution

Best Uses

- · Controls both goosegrass and crabgrass
- Improved bermudagrass control combination with triclopyr ester
- Controls Japanese stiltgrass in turf and ornamental areas

Attributes of Acclaim® Extra

- Systemic activity, rapidly absorbed through the foliage
- Tolerant on labeled turf species
- Broad-spectrum control of grassy weeds

Key Strengths

- Post-emergent annual and perennial grassy weed control
- For use in cool-season and zoysia turf, and around ornamentals





crabgrass

How To Apply

Goosegrass and Crabgrass Control

Acclaim Extra can be used to control both goosegrass and crabgrass in cool-season turf as well as zoysiagrass. Goosegrass in cool-season turf is a growing problem without many control options. Rates depend on the size of the weed.

ANNUAL GRASS WEED GROWTH STAGE	RATE (per acre)	RATE (per 1,000 sq ft)
1-leaf to 2-leaf ¹	3.5 fl oz	0.08 fl oz
Untillered ²	9.0 fl oz	0.21 fl oz
Untillered	13.0 fl oz	0.30 fl oz
1-tiller to 2-tiller	20.0 fl oz	0.46 fl oz
3-tiller to 4-tiller	28.0 fl oz³	0.64 fl oz
4-tiller to 5-tiller	39.0 fl oz³	0.90 fl oz

¹Apply this rate for use in multiple applications on bentgrass turf. ²Apply this rate only when tank mixing with pre-emergent herbicides. ³Do not apply more than 28.0 oz. of product to Kentucky bluegrass or zoysiagrass per application. Do not apply more than 39.0 oz. of product to tall fescue, fine fescue or annual and perennial ryegrass.



How To Apply (continued)

Bermudagrass Control

Bermudagrass is a major weed problem in tall fescue and zoysia turf, particularly in the transition zone of the United States. Not many options exist for controlling this tough perennial weed.

Tank mixes of Acclaim Extra and triclopyr ester may be used to effectively control bermudagrass in zoysiagrass and tall fescue when used in a treatment program. Use a tank mixture of Acclaim Extra at 20-28 fl. oz./acre and 16-32 oz. of triclopyr ester when bermudagrass becomes evident in zoysiagrass and tall fescue. Do not apply triclopyr ester to zoysiagrass unless some injury can be tolerated. For the best results, repeat applications every 4-6 weeks for up to 5 applications per season depending on your local conditions. Use the lower rates for uniform populations of bermudagrass intermixed with the desired turf, and higher rates where large bermudagrass patches are dominant in the turf stand. Do not apply Acclaim Extra at more than a total of 120 fl. oz./acre (2.75 fl. oz./1,000 sq. ft.) per growing season.

Japanese Stiltgrass Control

Japanese stiltgrass (Microstegium vimineum) is an annual grassy weed that is adapted to low-light conditions and was introduced into the U.S. in 1919. Common on roadsides, disturbed riparian areas and shaded areas with bare ground, it is considered an invasive species that outcompetes native plants. Japanese stiltgrass has invaded roadsides, parks, landscapes and recreation areas. Acclaim Extra is very effective on this weed and can be used on cool-season turf and around ornamentals, unlike many non-selective, post-emergent herbicides.

WEED GROWTH STAGE	RATE (per acre)	PINTS (per acre)
1-Tiller	16.0 fl oz	1.00
2-Tillers	20.0 fl oz	1.25
4-Tillers	28.0 fl oz	1.75
5-Tillers	39.0 fl oz	2.44

Spray Application For Spot Treatments

Add 40.0 fl. oz. of Acclaim Extra per 100 gal. of water, or 0.4 fl. oz./1.0 gal. of water and spray Japanese stiltgrass to point of spray deposit runoff.



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS CAREFULLY.



Bayer Crop Science L.P. Environmental Science Division, 2 TW Alexander Drive, Research Triangle Park, NC 27709, 1-800-331-2867, www.BackedByBayer.com Bayer, the Bayer Cross and Acclaim are registered trademarks of Bayer. Not all products are registered in all states. ©2015 Bayer CropScience LP. 15BPG-012-16

















Japanese stiltgrass