



For selective preemergence control of grass and broadleaf weeds in:

- Established Turf Grasses (excluding golf course putting greens) and lawns
- Landscape Ornamentals (including established perennials and wildflower plantings)

Active Ingredient
Prodiamine..... 0.42%
Other Ingredients..... 99.58%
Total..... 100.00%

¹(CAS # 2909-1212)

EPA Reg No.: 53883-297-94391

EPA Est. No.: 94391-KS-1

Guaranteed Analysis
Soluble Potash (K₂O) 7.00%

DERIVED FROM: Muriate of Potash.

KEEP OUT OF REACH OF CHILDREN CAUTION

(See the following section, Precautionary Statements, Hazards To Humans and Domestic Animals, for additional information.)

IF IN EYES:	FIRST AID Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor, for treatment advice.	
	If swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do by a poison control center or doctor. Do not give anything by mouth to an unconscious person.	
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.	
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.	

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 867-9050 for emergency medical treatment information.

Manufactured and Guaranteed By:
Mears Fertilizer, Inc.
629 North Industrial Road
El Dorado, KS 67042

NET WEIGHT: 50 Lbs. (22.68 Kgs.)

F 1958

Prodiamine Herbicide 0.42% Plus Turf Fertilizer 00-00-07

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS: This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

DIRECTIONS FOR USE:

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- * Established turf grasses (excluding golf course putting greens) and lawns,
- * Landscape ornamentals

- * Established perennials and wildflower plantings

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown for: (1) Sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Do not apply this product through any type of irrigation system.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

WEEDS CONTROLLED:

When used in accordance with this label this product will provide control of the following weeds:
Barnyardgrass, Bluegrass, Annual (Poa annua), Carpetweed, Chickweed - common, Chickweed, mouse ear (from seed), Crabgrass (large, smooth), Cowfootgrass, Cyngrass, Woolly, Foxtails, Annual, Goosegrass¹ Herbit, Itchgrass, Johnsongrass (from seed), Junglerice, Knotweed, Kochia, Lambsquarters - common, Lovegrass, Panicum (Texas, Fall, Browntop), Pigweed, Purslane - common, Pusley, Florida, Rescuegrass³ Shepherd's Purse², Signalgrass, Broadleaf, Speedwell, Persian, Sprangletop, Scurge, Prostrate, Violegress, Woodsorrel, Yellow (from seed)

¹In many areas a single application of 0.65lb. to 1.5 lb. active ingredient per acre (equal to 155 lbs. to 357 lbs./A of this product [or] equal to 3.6 to 6.2 lbs./10130 sq. ft. of this product) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 0.65 lb. to 1.0 lb. a.i. per acre (equal to 155 lbs. to 238 lbs./A of this product [or] equal to 3.6 to 5.5 lbs./1000 sq. ft. of this product) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above. ²Applications for this weed should be made in late summer, fall, or winter prior to germination. ³Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

APPLICATION DIRECTIONS:

Apply uniformly with suitable, calibrated application equipment.



ESTABLISHED TURF: This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map above for approximate crabgrass seed germination dates.

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

RATES OF APPLICATION: This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES: This product is for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate/Calendar Year of turf fertilizer by turf grass species ¹⁾			
Turf Species		Lbs. Product/A [or] ft.	Lbs. a.i/A
Creeping Bentgrass		155	3.6
Creeping Red Fescue		179	4.1
Bullhoggrass, Kentucky Bluegrass, Perennial Ryegrass		238	5.5
Bermudagrass ²⁾ , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia		357	8.2
			1.5

¹These are the maximum rates per calendar year by species limitations. ²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb. a.i./acre (equal to 119 lbs./A of this product [or] equal to 2.7 lbs./1000 sq. ft. of this product). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in springing situations.

* Do not apply more than 1.5 lbs. a.i. per calendar year per acre (equal to 357 lbs./A of this product [or] equal to 8.2 lbs./1000 sq. ft. of this product).

* Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates Table**.

WHEN TO APPLY AFTER OVERSEEDING TURF: Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not hatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the following table below for best overseeding/reseeding results.

Lbs. Product/Acre	Lbs Product/1000 sq. ft.	Lbs. a.i/A	MONTHS BEFORE OVERSEEDING		
			North	Transition	South
119	2.7	0.50	4	4	4
155	3.6	0.65	5	4	4
179	4.1	0.75	6	5	5
190	4.4	0.80	-	6	6
238	5.5	1.00	-	7	7
271	6.2	1.14	-	-	9
310	7.1	1.30	-	-	10
357	8.2	1.50	-	-	12

USE RATES AND WHEN TO APPLY: Apply in fall and/or spring, 0.65 to 1.5 lbs. active ingredient per acre, which is equal to 3.6 to 8.2 lbs. of this product per 1,000 sq. ft. of ornamental plantings; therefore 20 lbs of this product will treat 2,439 to 5,556 sq. ft. Use the higher rates of application for longer control periods. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rate of 1.5 lbs. a.i. per acre.

ESTABLISHED LANDSCAPE ORNAMENTALS: This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil. **NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS:** This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union.

THRIVE ENDURANCE is a trademark of Mears Fertilizer, Inc.



Sample Label

MFI # BP00750F042

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when it is uniformly applied to the soil surface, avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

[illegible]

*Do not use on food producing trees, vines or plants.
 **Not for use on container grown plants.

Label: PROD42 (New 12/26/23)

USE PRECAUTIONS AND RESTRICTIONS: The following precautions apply to the use of this product in turf grasses and lawns. (1) Application of this product may thin emerged plants in some areas. (2) Do not apply to overseeded turf within 60 days after seeding or until the second mowing, whichever is longer. Injury to annual bluegrass and newly overseeded grasses. (3) Do not apply to established seedlings in the second inch of soil, not touch plus soil. (4) Do not apply this product to putting greens or areas where dandelion, crabgrass, or other undesirable weeds are present. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens or areas where bentgrass, annual bluegrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species.

STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

CONDITIONS OF SALE AND WARRANTY (IMPORTANT: READ BEFORE USE)

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Meigs Fertilizer, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Means Fertilizer, Inc. makes no other warranties, expressed or implied, of merchantability or of fitness for a particular purpose, or otherwise, that extend beyond the statements made on this label. No agent of Means Fertilizer, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Means Fertilizer, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Weears Fertilizer, Inc's election, the replacement of product.

In Virginia, this lawn fertilizer contains phosphorous and is only for non-agricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf.

to Maryland application: The product being used in accordance with the recommended application rates established by the University of Maryland.

Black-Eyed Susan; Goldsturm
Santolina
Saxifrage; Purple Dome
Pincushion Flower
Stonecrop
Stonecrop
Stonecrop; Dragon's Blood
Stonecrop; Dragon's Blood
Mullein
Australian Boughery
Japanese Yew
Yew
Germander
Meadow Rue
American Abornifae
Star Jasmine
Catalpa Hemlock
Tulip species
Japanese Anemone
Sweet Viburnum
Japanese Snowball
Canary Island Viburnum
Arrowood Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Greater Penwinkle, Vinca
Common Penwinkle, Vinca
Coneflower
Old Fashioned Weigela
Spanish Bayonet, Yucca
Adam's Needle, Yucca

[illegible]

Lavender, Mustard
 Anise, Privet
 Chinese Privet, Variegata
 Japanese Privet
 Glossy Privet (Wickstea)
 Ivy, Juniper
 Big Blue Lily, Silver Mound,
 Evergreen Galt
 Red Flowering Currant
 Forsythia, Creeping
 Cardinal Flower, Indian Pink
 Japanese Honeysuckle
 Japanese Honeysuckle
 Burgundy
 Lonicera
 Magnolia species*
 Ice Plant
 Oleander
 Spiderbush
 Heavenly Bamboo
 Narcissus species*
 Oleander
 Olive*
 Moss Grass
 Osmanthus (Fatsia Holly), Gu
 Traveling African Daisy
 Sunwood
 Tree Holly
 Oriental Grass (Dwarf)
 Waveroad
 Fragrant Phlox (Red)
 Dragonhead, False Vio

[illegible][illegible][illegible]