

Bayer Solutions Quick Reference Guide - Fungicide Application Rates & Intervals

Go to www.BackedByBayer.com for complete product information
 Products may not be labeled in all U.S. states - check local registrations

	ARMADA® 50 WDG	CHIPCO® 26019	26GT®	BANOL®	BAYLETON® FLO	COMPASS® 50 WG	FIATA® SG	INTERFACE® SG	MIRAGE® SG¹	PROSTAR®	CHIPCO® SIGNATURE™	TARTAN® SG	CHIPCO TRITON® FLO
Golf													
Institutional Areas													
Lawn & Landscape²												no residential use	
Sod													
Sports Turf													
Pythium Control													
Plant Health												submitted	pending
Stressgard® Formulation Technology													
FRAC Code	3 + 11	2	2	28	3	11	33	2 + 11	3	7	33	3 + 11	3
USE RATES PER 1,000 SQ. FT. (OZ. OR FL. OZ.) / APPLICATION INTERVAL (DAYS)													
Disease	ARMADA 50 WDG	CHIPCO 26019	26GT	BANOL	BAYLETON FLO	COMPASS 50 WG	FIATA SG	INTERFACE SG	MIRAGE SG	PROSTAR	CHIPCO SIGNATURE	TARTAN SG	CHIPCO TRITON FLO
Algae and Cyanobacteria ●●													0.55-1.1/14-28
Anthraxnose ●●	0.6-1.5/14-28	4.0-8.0/special	4.0-8.0/special		1.0/30	0.1-0.25/14-21*		4.0-7.0/14	1.0-2.0/14-28		4.0-8.0/14*	1.0-2.0/14-28	0.41-1.1/14-28
Bacterial Decline (wilt) ●											8.0/7-14*		
Bentgrass Deadspot ●											4.0/14*		
Bipolaris Leaf Spot/Melting Out ●●	0.6-1.5/14-28	3.0-4.0/14-28	3.0-4.0/14-28			0.1-0.25/14-28		3.0-5.0/14-28				1.0-2.0/14-28	
Brown Patch ●	0.6-1.5/14-28	3.0-4.0/14-28	3.0-4.0/14-28		0.5-1.0/15-30*	0.1-0.25/14-21*		3.0-5.0/14-28	1.0-2.0/14-28	1.5-3.0/14-21*		1.0-2.0/14-28	0.41-1.1/14-28
Brown Ring (Waitea) Patch ●									1.0-2.0/14-28	2.2-4.5/30			0.55-1.1/14-28
Copper Spot ●●					0.5-1.0/15-30								
Curvularia Blight ●●		4.0-8.0/14	4.0-8.0/14					3.0-7.0/14					
Dollar Spot ●●	0.6-1.5/14-28	3.0-4.0/14-21*	3.0-4.0/14-21*		0.25-1.0/14-30			3.0-5.0/14-21*	1.0-2.0/14-28			1.0-2.0/14-28	0.28-1.1/14-28
Fairy Ring ●●	1.2-1.5/14-28				1-1.9/14-21				1.0-2.0/28	2.2-4.5/21-30		2.0/special	0.55-1.1/28s
Fusarium Blight ●		8.0/28	8.0/28		1.0-1.9/30			7.0/28					
Gray Leaf Spot ●●	0.6-1.5/14-28				0.5-1.0/14	0.15-0.25/14-21*			1.0-2.0/14-28			1.0-2.0/14-28	
Large/Zoysia Patch ●	1.2-1.5/28	4.0/14-21	4.0/14-21		1.0-1.9/special			4.0/14-21	1.0-2.0/28	2.2/30			0.55-1.1/14-28
Leaf and Sheath Spot ●●										2.2-4.5/14-21			
Microdochium (Fusarium) Patch ●●	1.2-1.5/special	4.0-8.0/14-21*	4.0-8.0/14-21*		1.0-1.9/60-90	0.2-0.25/special*		4.0-7.0/14-21*	1.0-2.0/10-14			2.0/special	0.28-1.1/10-14
Necrotic Ring Spot ●	1.2-1.5/28	8.0/28	8.0/28					7.0/28	2.0/28				0.28-0.55/28
Pink Patch ●●	0.6-1.5/14-28					0.15-0.25/14-21*		3.0-4.0/14	1.0-2.0/14-28	1.5/21-28		1.0-2.0/14-28	0.55-1.1/14-28
Powdery Mildew ●●					0.5-1.0/15-30								
Pythium Blight ●●				1.3-4.0/7-21*			4.4-15.0/14-28*				4.0-8.0/14-21		
Pythium Damping Off ●●				1.3-4.0/7-21*									
Pythium Root Dysfunction ●				2.0/special *							4.0/special*		
Pythium Root Rot ●●				1.3-4.0/7-21*			4.4-15.0/14-28*				4.0-8.0/14-21		
Rapid Blight ●	0.6-1.5/14-28*					0.15-0.25/14-21*		3.0-5.0/14-28					
Red Thread ●	0.6-1.5/14-28	4.0/14	4.0/14		0.5-1.0/15-30	0.15-0.25/14-21*		3.0-4.0/14	1.0-2.0/14-28	1.5/21-28		1.0-2.0/14-28	0.41-1.1/14-28
Rust ●●	0.6-1.5/14-28				0.5-1.0/15-30	0.1-0.25/14-21*		3.0-5.0/14-28	1.0-2.0/14-28			1.0-2.0/14-28	0.28-0.55/14-28
Snow Mold (Gray) ●		4.0-8.0/special	4.0-8.0/special		1.9/special*			4.0-7.0/special	1.0-2.0/28	3.0-4.5/special			0.28-1.1/14-28
Snow Mold (Pink) ●	1.2-1.5/special	4.0-8.0/special	4.0-8.0/special		1.0-1.9/special	0.25/special*		4.0-7.0/special	1.0-2.0/28			2.0/special	
Southern Blight ●●	0.6-1.5/14				0.5-1.9/14-28					1.5/21-28		1.0-2.0/14	
Spring Dead Spot ●									2.0/28				
Stripe Smut ●	0.6/special				1.0/special							1.0/special	
Summer Decline/Stress ●●		2.0-4.0/special*	2.0-4.0/special*					3.0-7.0/14*			4.0-8.0/14*	1.5-2.0/28	0.55-1.1/14-28
Summer Patch ●	1.2-1.5/21-28				1.0-1.9/30	0.2-0.25/21-28*			1.0-2.0/14-28			2.0/21-28*	0.55-1.1/14-28
Take-All Patch ●	1.2-1.5/28				1.0-1.9/21-28*				1.0-2.0/14-28				0.55-1.1/14-28
Take-All Root Rot/Decline ●					1.0-1.9/21-28				2.0/28				
Yellow Patch ●●									1.0-2.0/21-28	1.5/21-28			0.55-1.1/14-28
Yellow Tuft ●											4.0-8.0/14-21		
Use rates²	0.6-1.5 oz	2.0-8.0 fl oz	2.0-8.0 fl oz	1.3-4.0 fl oz	0.25-1.9 fl oz	0.1-0.25 oz	4.4-15.0 fl oz	2.0-7.0 fl oz	1.0-2.0 fl oz	1.5-4.5 oz	4.0-8.0 oz	1.0-2.0 fl oz	0.28-1.1 fl oz
Max use/season (oz)²	4.75 oz	35.0 fl oz	35.0 fl oz	12.0 fl oz	3.8 fl oz	0.8 oz		30.0 fl oz	6.5 fl oz (3.0 in NY)	4.5 oz²		7.9 fl oz	3.3 fl oz
Max app/season²		6	6					6	6.0 (3.0 in NY)				
Other restrictions²					≤1.4 fl oz for L & L4				3 x 1.0 fl oz in NY				No NY Use

● = cool season
 ●● = warm season

All rates, diseases and intervals should be verified on the approved state label associated with product packaging. Not all products, rates and diseases are approved in all states. Please consult labels prior to applying.
 *See label, 2(ee) supplemental or Bayer Solutions Guide for additional application or tank-mix instructions. *Mirage SG may not be registered for use in CA & NY until later in 2014. *Rate info for most products shown for Golf use, check label for specific Lawn and Landscape instructions. *Additional spot treatments can be made for fairy ring. *Bayleton has a max rate of 1.4 fl. oz./1,000 sq. ft. per application on Lawn and Landscape. ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS CAREFULLY. ©2015 Bayer CropScience LP.

