



# Proxy<sup>®</sup>

**Active Ingredient:** ethephon  
**Formulation:** flowable  
**Signal Word:** danger  
**Activity:** plant growth regulator

## Best Uses

- Suppresses the formation of seedheads of various plants including *Poa annua* and white clover
- Suppress the growth of labeled cool-season grasses

## Attributes Of Proxy<sup>®</sup>

- Foliarly absorbed, and is most effective on actively growing healthy turf
- Rainfast within 2 hours
- Promotes turf that is denser, tighter-knit and more playable
- Tank-mix with Primo MAXX for seedhead suppression and improved turf quality

## Key Strengths

- *Poa annua* seedhead suppression in combination with Primo MAXX<sup>®</sup>
- Regulates growth of desirable grasses in cool-season turf, especially roughs and tall grass



*Poa annua*

## How To Apply

- For best results, apply in sufficient volume of water to provide uniform coverage
- For maximum performance, delay mowing until the day after application
- Make the initial application prior to the emergence of new seedheads
- A period of 2-3 weeks after application is required for maximum performance
- Repeat applications may be made to predominately *Poa annua* or white clover sites as needed, but wait two weeks after the previous application
- Follow other general recommendations and precautions on the label



# Proxy®

APPLICATION	SITES	RATE (per 1,000 sq ft)	SPRAY VOL. (per 1,000 sq ft)
<b><i>Poa annua</i> and white clover seedhead suppression:</b>  <b>Reapplication interval: 2 weeks or greater for all labeled grasses</b>	<b>Commercial turfgrasses including bentgrass, Kentucky bluegrass, perennial ryegrass, tall and fine fescue and bermudagrass</b>  <b>Golf course turf including greens, tees, fairways and roughs</b>	<b>5.0 fl oz</b>	<b>1.0 - 2.0 gal</b>
<b>Turfgrass growth regulation reapplication intervals:</b> <ul style="list-style-type: none"> <li>• Kentucky bluegrass - 7 weeks</li> <li>• Perennial ryegrass - 7 weeks</li> <li>• Tall and fine fescue - 4 weeks</li> <li>• Bentgrass - 4 weeks</li> </ul>	<b>Golf course turf including greens, tees, fairways and roughs</b>  <b>Commercial turfgrasses including bentgrass, Kentucky bluegrass, perennial ryegrass, tall and fine fescue</b>	<b>5.0 fl oz</b>	<b>1.0 - 2.0 gal</b>



Close-up of plugs from an untreated and treated area respectively. Seedhead suppression results in healthier plants and improved turf quality. Photo: Mike Hirvela, Bayer CropScience.



Proliferation of annual bluegrass seedhead in an untreated area on a golf course putting green. Photo: Mike Hirvela, Bayer CropScience.

BackedByBayer.com | @BayerGolf | | Available on iTunes and Google play: BackedByBayer



**ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS CAREFULLY.**

Bayer CropScience LP, Environmental Science Division. 2 TW Alexander Drive, Research Triangle Park, NC 27709. 1-800-331-2867. [www.BackedByBayer.com](http://www.BackedByBayer.com). Bayer, the Bayer Cross and Proxy are registered trademarks of Bayer. Primo MAXX® is a registered trademark of Syngenta. Not all products are registered in all states. ©2015 Bayer CropScience LP. 14BPG-052-P4

Backed by  
**BAYER**