

Active Ingredients: indaziflam Formulation: flowable WSSA Group: 29 Signal Word: no signal word required

Best Uses

- Pre-emergent herbicide controls the most troublesome grassy and broadleaf weeds, as well as annual sedges
- Extended residual control of annual bluegrass (*Poa annua*), goosegrass, crabgrass and 75 more weeds

Attributes Of Specticle[®] FLO

- Controls weeds by inhibiting cellulose biosynthesis

 its primary site of action is the newly developing root of a germinated weed seedling
- No stains, no odor formulation
- Mixable with post-emergent herbicides
- Newer mode-of-action controls weeds known to develop resistance to other herbicides
- Works in turf, landscapes and hardscapes

How To Apply

- Apply to turf before weed seeds germinate
- Use as a broadcast treatment or directed spray on bare ground or mulched areas around labeled ornamental plants and grasses
- Adequate soil moisture is needed to activate Specticle FLO and provide maximum control
- If weeds are present, tank-mix Specticle FLO with selective, post-emergent herbicides like Revolver[®], Celsius[®] WG or Tribute[®] Total
- To control emerged weeds in plant beds, dormant turf, cart paths or parking lots, use in conjunction with a non-selective herbicide

Key Strengths

- Broad spectrum pre-emergent weed control
- Long residual
- Poa annua control
- Low use rates



Poa annua

Eleusine indica

Digitaria sanguinalis



Use Rates For Single Application Of Specticle FLO

Target Weed (single application)	Specticle FLO Use Rate (fl oz per acre)
Crabgrass Goosegrass Annual bluegrass Broadleaf weeds	6.0 - 10.0
Annual sedge Kyllinga	10.0

Use Rates For Split Application Of Specticle FLO

Target Weed	Initial Application Use Rate (fl oz per acre)	Follow-up Application (within 45 - 90 days) Use Rate (fl oz per acre)
Crabgrass Goosegrass Annual bluegrass	4.5 - 9.0	3.0 - 9.0
Broadleaf weeds	6.0 - 9.0	3.0 - 9.0
Annual sedge Kyllinga	9.0	6.0 - 9.0

Use Rates For Multiple Applications Of Specticle FLO

Target Weed	Initial Application Use Rate (fl oz per acre)	Second Application (within 30 - 45 days) Use Rate (fl oz per acre)	Third Application (within 30 - 45 days) Use Rate (fl oz per acre)
Crabgrass Goosegrass Annual bluegrass*	3.0 - 4.5	3.0 - 4.5	3.0 - 4.5

*Specticle FLO 2(ee) allows these use rates on annual bluegrass in the following states: AL, AZ, AR, CO, FL, GÁ, HI, IN, IL, KS, LÁ, MD, MS, MO, NC, OH, OK, SC, TN, TX, UT, VA, WV.



goosegrass



large crabgrass



spotted spurge



doveweed

BackedByBayer.com





ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS CAREFULLY. Bayer CropScience LP, Environmental Science Division. 2 TW Alexander Drive, Research Triangle Park, NC 27709. 1-800-331-2867. www.BackedByBayer.com. Bayer, the Bayer Cross, Celsius, Revolver, Specticle and Tribute are registered trademarks of Bayer. Not all products are registered in all states. @2015 Bayer CropScience LP. 14BP6-052-P10