



FLUAZINAM 40SC SELECT™

FUNGICIDE

Broad Spectrum Turf Fungicide



ACTIVE INGREDIENT

Fluazinam

% ACTIVE INGREDIENT

40%

FORMULATION

Soluble concentrate

CHEMICAL FAMILY

Quinone inside Inhibitors
(Qil); Group 29 Fungicide

BRAND ALTERNATIVE

Secure®

SIGNAL WORD

Warning

PACKAGING INFORMATION

2 x 2.5 gal

LABELED USES

Golf course tees, greens,
fairways and roughs

Fluazinam 40SC Select™ is a broad spectrum, multi-site fungicide to control diseases in turf grasses on golf courses only. The active ingredient, fluazinam, is the only registered fungicide for turf in FRAC group 29. It has no known resistance and as a multi-site contact, it has a very low risk of developing resistance.

Fluazinam 40SC Select™, when partnered with Chlorothalonil-based fungicides, provides season-long protection against the toughest turf diseases without interruptions.

Fluazinam 40SC Select™ not only provides superior protection against dollar spot, but also controls 10 additional turf diseases. It even controls dollar spot that is resistant to other chemistries.

- Unique multi-site contact active ingredient, reduces resistance management
- Excellent control of dollar spot and 10 additional turf diseases
- Perfect tank mix partner with systemics for inside out protection
- Controls dollar spot that is resistant to other chemistries

IMPORTANT:
ALWAYS READ AND FOLLOW
LABEL INSTRUCTIONS.


PRIME SOURCE

©2017 Prime Source, LLC. Important: Always read and follow label instructions for use. Secure is a registered trademark of Syngenta Crop Group.

Prime Source, LLC
10025 US 264 Alternate
Middlesex, NC 27557
www.prime-sourcellc.com

FLUAZINAM 40SC SELECT™

FUNGICIDE

Use and Application Information

Apply Fluazinam 40SC Select™ in sufficient water to obtain adequate uniform coverage of the turf. Apply in a minimum spray volume of 40 gal of water per acre on fairways and 90 gal of water per acre on tees and greens.

- Apply at 0.5 fl. oz. per 1,000 sq. ft. (6 acres per gallon) every 14 days.
- Apply after mowing with sufficient water to uniformly cover the turf and allow area to dry completely before irrigation.
- Refer to the product label for additional information about this product and tank mixing.
- Not for use on home lawns or any residential turf.

Turfgrass Disease Control

- Use Fluazinam 40SC Select™ in a preventative disease control program.
- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed area to completely dry before mowing.
- For control of foliar diseases, allow sprayed area to completely dry before irrigation.
- For optimum turf quality and disease control, use Fluazinam 40SC Select™ in conjunction with turf management practices that promote good plant health and optimum disease control.
- Evaluate spray additives prior to use. Label directions are based on data obtained with no additives.
- Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.



Restrictions

- DO NOT exceed 258 fl oz of Fluazinam 40SC Select™ (8.4 pounds active ingredient) per acre, per year.
- DO NOT apply more than twelve (12) applications at 21.5 fl oz per acre (0.5 fl oz/1000 sq ft), per year

DISEASES CONTROLLED INCLUDE:

Algal scum
Anthracnose
Brown blight
Brown patch
Dollar spot
Gray snow mold
Large patch
Leaf spot
Melting out
Pink patch
Pink snow mold
Red thread
Stem rust



Prime Source, LLC
10025 US 264 Alternate
Middlesex, NC 27557
www.prime-sourcellc.com

Download a
specimen label at
Prime-Sourcellc.com